

PROTEASE

Transformation's most
therapeutic systemic
proteolytic formula

This proprietary blend of highly active, GI stable proteolytic enzymes has been combined to promote circulation, a strong healthy immune system, reduced inflammation, and timely detoxification.

Product Highlights

- Endo/exo peptidases break the inner/terminal bonds of amino acid chains for more efficient hydrolysis of proteins
- Protease blend (including bromelain plant enzymes) for reducing inflammation
- More than 400,000 HUT for the highest proteolytic activity available (600,000 PU = 51,000 HUT)
- 19 SAPU units from Protease 3.0
- Approx. 2,400 FU breaks down fibrin and clots and promotes healthy blood flow
- Calcium improves tolerance on an empty stomach

| SUPPLEMENT FACTS | |
|--|---------------|
| Serving Size 1 Capsule | |
| Amount Per Serving | % Daily Value |
| Tzyme™ Protease Blend (peptidases, bromelain) (355,020 HUT + 19 SAPU) (600,000 PU) | 492 mg † |
| † Daily Value not established | |

Other Ingredients: Vegetable Capsule (Hydroxypropylmethylcellulose, Water), Calcium Citrate

Clinical Applications

- Inflammatory disorders (GI, joints, skin, respiratory system, and organs)
- Support and prevention for cancer
- Immune system imbalances
- Hypertension
- Chronic Fatigue / Fibromyalgia
- Bacterial / Viral / Fungal Infections
- Hepatitis
- Kidney Disorders / Renal Insufficiency
- Skin disorders
- Chronic respiratory issues
- All hormone imbalances
- Auto-immune disorders
- Support for complications associated with autism
- Support for complications associated with diabetes
- Muscle strains, soreness, injuries, and surgeries

For Your Information

- First choice when patient has been diagnosed with a condition (this is the "therapeutic" strength blend)
- It is better to take small doses of protease frequently throughout the day rather than large doses once or twice a day (protease has a half-life of approximately 3-4 hours; the goal is to keep the protease activity constant in the blood stream for therapeutic benefits)
- Compares with Nattokinase and Serra-peptidase
- May be given to children if condition warrants; may be given to pets
- Caution with patients on prescription blood thinning drugs (give protease formulas 3-4 hours away from Rx dose)
- May cause discomfort for individuals with stomach ulcers as protease will debride necrotic tissue and promote healing
- Discontinue taking Protease 24-48 hours prior to surgery and resume 24 hours post-surgery

Dosage

- Maintenance dose: 1 capsule 3 x day (rise, midday, rest) on an empty stomach
- General therapy dose: 2 capsules 3 x day between meals (approximately 1 hour before or 2 hours after meals)
- Therapeutic dose: 2-3 capsules 4-5 x day, or every 3 hours (goal is to keep high levels of activity in the blood stream at all times; "between meals" becomes difficult, so 30-60 minutes before or after meals is acceptable)
- **MAY BE TAKEN WITH FOOD IF UNABLE TO TOLERATE BETWEEN MEALS**
- Sometimes it is suggested 1 capsule may be taken with meals to enhance the digestion of proteins (diabetes, heartburn, acid reflux, gout, autism, high protein diet)
- For those who have difficulty swallowing pills, the capsules may be pulled apart and mixed in a small amount of tepid water or liquid and consumed immediately
- Topical application: open capsule and make a paste with a small amount of water, then apply to insect bite, fungal rash, mouth sores, inflamed gums, etc